

# ANNALES DE L'I. H. P., SECTION A

L. MANGIAROTTI

M. MODUGNO

## **Some results on the calculus of variations on jet spaces**

*Annales de l'I. H. P., section A*, tome 39, n° 1 (1983), p. 29-43.

[http://www.numdam.org/item?id=AIHPA\\_1983\\_\\_39\\_1\\_29\\_0](http://www.numdam.org/item?id=AIHPA_1983__39_1_29_0)

© Gauthier-Villars, 1983, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A », implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## Some results on the calculus of variations on jet spaces (\*)

by

L. MANGIAROTTI (\*\*) and M. MODUGNO (\*\*\*)

---

**ABSTRACT.** — The basic object is a fibered manifold  $p: E \rightarrow M$  and the framework is that of jet spaces. Given a Lagrangian form  $\Lambda$  on  $JE$ , we work with the space  $\{\Lambda\}$  of variational forms associated to  $\Lambda$ . It is this space which is important in the calculus of variations. We study a new operator  $\mathcal{E}_\Lambda$  (defined only on  $\text{Ker } \vartheta \subset TJE$  where  $\vartheta$  is the fundamental 1-form) canonically associated to  $\{\Lambda\}$ . This operator is well suited for studying critical sections and functorial properties. The so called Euler-Lagrange operator  $E_\Lambda$  appears as an extension of  $\mathcal{E}_\Lambda$ . Variational symmetries are introduced as morphisms of a category whose objects are the variational forms  $\{\Lambda\}$ . The uniqueness of the Poincaré-Cartan form  $\Theta_\Lambda$  is proved under certain circumstances. Various interesting relations between  $\Lambda$ ,  $E_\Lambda$  and  $\Theta_\Lambda$  are investigated.

**RÉSUMÉ.** — L'objet fondamental est une variété avec un espace fibré  $p: E \rightarrow M$  et le cadre est celui des espaces de jets. Étant donnée une forme Lagrangienne  $\Lambda$  sur  $JE$ , on considère l'espace  $\{\Lambda\}$  des formes variationnelles associées à  $\Lambda$ , qui est l'espace important pour le calcul des variations. On étudie un nouvel opérateur  $\mathcal{E}_\Lambda$  (défini seulement sur  $\text{Ker } \theta \subset TJE$ , où  $\theta$  est la 1-forme fondamentale), canoniquement associé à  $\{\Lambda\}$ . Cet opérateur est bien adapté à l'étude des sections critiques et des propriétés fonctorielles. L'opérateur d'Euler-Lagrange  $E_\Lambda$  apparaît comme extension de  $\mathcal{E}_\Lambda$ . Les symétries variationnelles sont introduites

---

(\*) Work performed under the auspices of C. N. R. (Gruppo Nazionale per la Fisica Matematica).

(\*\*) Permanent address: Istituto di Matematica, Università, Camerino (MC).

(\*\*\*) Permanent address: Istituto di Matematica Applicata, Università, Firenze.

comme morphismes d'une catégorie dont les objets sont les formes variationnelles de  $\{\Lambda\}$ . On montre l'unicité de la forme  $\Theta_\Lambda$  de Poincaré-Cartan dans certaines circonstances. On étudie quelques relations intéressantes entre  $\Lambda$ ,  $E_\Lambda$  et  $\Theta_\Lambda$ .

---

## INTRODUCTION

In recent years much work has been done on the calculus of variations on jet spaces [2] [3] [4] [6] [7] [9] [10] [11] and [12]. Following this stream, the present paper emphasizes both the geometrical basis of the theory as well as its functorial nature. Moreover, it contains novelties in the treatment as well as new results.

The basic geometric object is a fibered manifold  $p : E \rightarrow M$  and the basic spaces are the first two jet spaces, i. e.  $JE$  and  $J^2E$ . In Section 1 we recall the two main facts about the geometry of jet spaces, namely the affine structures and the fundamental 1-form  $\vartheta$ . Then the subject of infinitesimal contact transformations is developed in a new way. These transformations are important because a « variational problem » is « infinitesimally functorial » with respect to them. A deeper analysis at all orders of jet spaces will appear in a related paper [8].

In Section 2 we introduce Lagrangian forms  $\Lambda$  on  $JE$ , i. e. horizontal forms with respect to the canonical projection  $JE \rightarrow M$  (in physical terms, these are Lagrangian densities). Affine Lagrangian forms are studied in detail. Then we define the space  $\{\Lambda\}$  of variational forms associated with a Lagrangian form  $\Lambda$  (already used in [10] in the context of affine bundles). It is  $\{\Lambda\}$  which is important in the Calculus of Variations. Various properties of  $\{\Lambda\}$  in relation with automorphisms  $\varphi : E \rightarrow E$  (for brevity we consider only automorphisms of  $E$  though many properties hold in a more general context) as well as infinitesimal contact transformations are demonstrated. Moreover, the uniqueness of the Poincaré-Cartan form is proved under certain circumstances.

A main result in Section 3 is the introduction of a new operator  $\mathcal{E}_\Lambda$  (defined only on  $\text{Ker } \vartheta \subset TJE$ ) canonically associated to the set  $\{\Lambda\}$ . This operator is well suited for studying critical sections and functorial properties. The classical Euler-Lagrange operator  $E_\Lambda$  is an extension of  $\mathcal{E}_\Lambda$ . Various interesting relations between  $\Lambda$ ,  $E_\Lambda$  and  $\Theta_\Lambda$  are investigated. Variational symmetries are introduced as morphisms of a category whose objects are the variational forms  $\{\Lambda\}$ . However, this is only a first insight into such an intricate question [2] [4] [6] and [11]. The analysis of the subject, also with respect to Noether's theorem, will be pursued in a future work.

In Section 4 we treat of critical sections. The functorial nature of a variational problem is exhibited in a very direct way. A new characterization of critical sections in terms of  $\mathcal{E}_\lambda$  is given.

### 1. NOTATION AND PRELIMINARY RESULTS

a) All manifolds and maps will be  $C^\infty$ . The notation used is that of modern differential geometry. The basic object is a fibered manifold  $p : E \rightarrow M$  (i. e.  $p$  is a surjective submersion),  $\dim M = m, \dim E = m + l$ . Fibered charts are denoted by  $(x^\alpha, y^i), 1 \leq \alpha \leq m, 1 \leq i \leq l$ . Induced fibered charts on the 1-jet space  $JE$  are denoted by  $(x^\alpha, y^i, y_\alpha^i)$ . The canonical projection  $JE \rightarrow E$  is denoted by  $p_E$ . Then  $p_M = p \circ p_E$  is the canonical projection  $JE \rightarrow M$ . If  $s$  is a (local) section of  $E$ , then  $js : M \rightarrow JE$  is the 1-jet prolongation of  $s$ . If  $\varphi : E \rightarrow E$  is an automorphism of  $E$  (over the diffeomorphism  $\varphi_M : M \rightarrow M$ ), then  $J\varphi : JE \rightarrow JE$  is the 1-jet prolongation of  $\varphi$ . Recall that  $J\varphi$  is characterized by  $J\varphi \circ js = j(\varphi_*s) \circ \varphi_M$  for any section  $s$  of  $E, \varphi_*s = \varphi \circ s \circ \varphi_M^{-1}$ .

There are two basic properties concerning  $JE$ . First,  $JE$  is an affine bundle over  $E$  with vector bundle the tensor product bundle  $T^*M \otimes_{\mathbb{R}} VE$  [3].

Here  $T$  and  $V$  denote the tangent and vertical functors, respectively ( $VE = \text{Ker } Tp \subset TE$ ). As a rule, obvious pull-backs will be omitted.

Second, there is a (first order) canonical inclusion over  $JE \rightarrow E$ , namely

$$(1.1) \quad c : JE \times_M TM \rightarrow TE$$

characterized by  $c \circ (js, u) = Tjs \circ u$  for any section  $s : M \rightarrow E$  and any vector field  $u : M \rightarrow TM$ . It induces the canonical projection

$$(1.2) \quad \mathcal{G} : JE \times_E TE \rightarrow VE,$$

the so called structure 1-form (see [2] and [3]), whose local expression is

$$(1.3) \quad (x^\alpha, y^i, y_\alpha^i; \dot{y}^i) \circ \mathcal{G} = (x^\alpha, y^i, y_\alpha^i; \dot{y}^i - y_\alpha^i \dot{x}^\alpha).$$

By means of the tangent map  $Tp_E : TJE \rightarrow TE, \mathcal{G}$  can also be considered as a linear morphism  $\mathcal{G} : TJE \rightarrow VE$  over  $p_E : JE \rightarrow E$ . Hence  $\mathcal{G}$  is a 1-form on  $JE$  valued over  $VE$  whose local expression is

$$(1.4) \quad \mathcal{G}^i = dy^i - y_\alpha^i dx^\alpha.$$

In the sequel, the subbundle  $\Delta = \text{Ker } \mathcal{G} \subset TJE$  will be used. A local basis of  $\Delta$  is given by  $(\partial_\alpha + y_\alpha^i \partial_i, \partial_i^?)$ . Here  $(\partial_\alpha, \partial_i, \partial_i^?)$  are the local fields

associated to the fibered charts  $(x^\alpha, y^i, y_\alpha^i)$ . Let us note that  $\mathfrak{g}$  (and hence  $\Delta$ ) is invariant with respect the automorphisms  $\varphi : E \rightarrow E$ , i. e.

$$(1.5) \quad \mathfrak{g} \circ \text{TJ}\varphi = \text{V}\varphi \circ \mathfrak{g}.$$

In the usual pull-back notation, (1.5) is written as  $\varphi^*\mathfrak{g} = \mathfrak{g}$ .

All that we have said holds for higher order jet spaces [3] [4] and [8]. Nevertheless we need only to consider the 2-jet space  $J^2E$  (because the Euler-Lagrange equation is of the second order). Induced fibered charts on  $J^2E$  are denoted by  $(x^\alpha, y^i, y_\alpha^i, y_{\alpha\beta}^i)$ . Note that there is a canonical injection  $J^2E \subset J(\text{JE})$ .

The 2-jet space  $J^2E$  is an affine bundle over  $\text{JE}$  with vector bundle  $(\bigvee_2 \text{T}^*\text{M}) \otimes_{\text{JE}} \text{VE}$ , where  $\bigvee_2$  denotes the second order symmetric tensor product.

From the second order canonical inclusion

$$(1.6) \quad c : J^2E \times_{\text{M}} \text{TM} \rightarrow \Delta \subset \text{TJE},$$

we get the (second order) structure 1-form on  $J^2E$  which can be considered as a linear morphism  $\text{TJ}^2E \rightarrow \text{VJE}$  over the canonical projection  $J^2E \rightarrow E$  (here  $\text{VJE} = \text{Ker Tp}_{\text{M}} \subset \text{TJE}$ ). Its local expression is

$$(1.7) \quad \mathfrak{g}^i = dy^i - y_\alpha^i dx^\alpha, \quad \mathfrak{g}_\alpha^i = dy_\alpha^i - y_{\alpha\beta}^i dx^\beta.$$

b) There are distinguished vector fields on  $\text{JE}$ , namely the so called infinitesimal contact transformations (in short : i. c. t.) [2] [4]. They are important in the calculus of variations, essentially because a « variational problem » is « infinitesimally functorial » with respect to them. We give here a new treatment of these vector fields (compare with [2] and [4]). A comprehensive analysis of the subject, involving jet spaces of any order, will appear in [8].

DEFINITION. — A vector field  $u$  on  $\text{JE}$  is an i. c. t. (of first order) iff  $L_u \Delta \subset \Delta$  (here  $L_u$  is the Lie derivative with respect to  $u$ ).

Let  $u = u^\alpha \partial_\alpha + u^i \partial_i + u_\alpha^i \partial_\alpha^i$  be the local expression of  $u$ . Then it is easily proved that  $u$  is an i. c. t. iff

$$(1.8) \quad u_\alpha^i = \partial_\alpha u^i + y_\alpha^j \partial_j u^i - y_\beta^j (\partial_\alpha u^\beta + y_\alpha^j \partial_j u^\beta), \quad \partial_\alpha^j u^i - y_\beta^j \partial_\alpha^i u^\beta = 0.$$

There is an interesting way of characterizing the i. c. t. which is based on the existence of a canonical projection  $r : \text{JTE} \rightarrow \text{TJE}$  [8]. Let us remark preliminarily that  $\text{JTE}$  is an affine bundle over the fibered product  $\text{JE} \times_{\text{M}} \text{JTM}$  with vector bundle (the pull-back over  $\text{JE} \times_{\text{M}} \text{JTM}$  of)  $\text{VJE}$ . Also  $\text{TJE}$  is an affine bundle over  $\text{JE} \times_{\text{M}} \text{TM}$  with vector bundle (the pull-back over  $\text{JE} \times_{\text{M}} \text{TM}$  of)  $\text{VJE}$ .

**PROPOSITION.** — There is an unique affine morphism  $r: JTE \rightarrow TJE$  over the canonical projection  $JE \times_M JTM \rightarrow JE \times_M TM$  such that

- i)  $r \circ JTs = Tjs$  for any section  $s: M \rightarrow E$ ,
  - ii) its fiber derivative  $Dr$  restricts to the identity over the fibers.
- The local expression of  $r$  is

$$(1.9) \quad (x^\alpha, y^i, y_\alpha^i, \dot{x}^\alpha, \dot{y}^i, \dot{y}_\alpha^i) \circ r = (x^\alpha, y^i, y_\alpha^i, \dot{x}^\alpha, \dot{y}^i, \dot{y}_\alpha^i - y_\beta^i \dot{x}^\beta).$$

**PROPOSITION.** — A vector field  $u$  on  $JE$  is an i. c. t. iff we have  $u = r \circ J(Tp_E \circ u)$ . If  $u_E$  is a vector field on  $E$ , then  $u = r \circ Ju_E$  is the unique i. c. t. projectable over  $u_E$  (1-jet prolongation of  $u_E$ ). Moreover, in the particular case when  $u_E$  is projectable over  $M$ ,  $u$  is the infinitesimal generator of  $JF_t$ , where  $F_t$  is the flux of  $u_E$ .

If  $u_E = u^\alpha \partial_\alpha + u^i \partial_i$  is the local expression of  $u_E$ , then the local expression of the 1-jet prolongation of  $u_E$  is

$$(1.10) \quad u = u^\alpha \partial_\alpha + u^i \partial_i + u_\alpha^i \partial_i^\alpha,$$

where  $u_\alpha^i$  is given by (1.8).

It is obvious how the previous definition works on  $J^2E$ . If

$$u = u^\alpha \partial_\alpha + u^i \partial_i + u_\alpha^i \partial_i^\alpha + u_{\alpha\beta}^i \partial_i^{\alpha\beta}$$

is the local expression of a vector field on  $J^2E$ , it follows from (1.7) that  $u$  is a second order i. c. t. iff

$$(1.11) \quad \begin{cases} u_\alpha^i = \partial_\alpha u^i + y_\alpha^j \partial_j u^i + y_{\alpha\beta}^j \partial_j^\beta u^i - y_\beta^i (\partial_\alpha u^\beta + y_\alpha^j \partial_j u^\beta + y_{\alpha\gamma}^j \partial_j^\gamma u^\beta), \\ u_{\alpha\beta}^i = \partial_\alpha u_\beta^i + y_\alpha^j \partial_j u_\beta^i + y_{\alpha\gamma}^j \partial_j^\gamma u_\beta^i - y_{\beta\gamma}^i (\partial_\alpha u^\gamma + y_\alpha^j \partial_j u^\gamma + y_{\alpha\lambda}^j \partial_j^\lambda u^\gamma), \\ \partial_j^{\alpha\beta} u^i - y_\gamma^i \partial_j^\beta u^\gamma = 0, \quad \partial_j^{\alpha\beta} u_\gamma^i - y_{\gamma\lambda}^i \partial_j^{\alpha\beta} u^\lambda = 0. \end{cases}$$

It follows easily from (1.8) and (1.11) that if  $u$  is projectable over a vector field  $u_E$  on  $E$  and also over a vector field on  $JE$ , then this projection is the 1-jet prolongation of  $u_E$ . As in the last proposition, it is clear that (1.11) gives us the local expression of the 2-jet prolongation of a vector field  $u_E$  on  $E$ .

## 2. VARIATIONAL FORMS

a) Let us recall that  $p_M$  denotes the canonical projection  $JE \rightarrow M$ . A  $p_M$ -horizontal  $m$ -form on  $JE$  is called a Lagrangian form on  $JE$ . An  $m$ -form  $\Lambda$  on  $JE$  is a Lagrangian form iff its local expression is

$$(2.1) \quad \Lambda = \mathcal{L}\omega, \quad \omega = dx^1 \wedge \dots \wedge dx^m,$$

where  $\mathcal{L}$  is a local function on  $JE$ .

If the manifold  $M$  is orientable, then the choice of a volume form on  $M$  induces a bijection between Lagrangian forms and functions on  $JE$ . In physical terms,  $\Lambda$  is a Lagrangian density while  $\mathcal{L}$  is a Lagrangian.

Let us make some remarks about Lagrangian forms of the affine type. Clearly,  $\Lambda$  is an affine map (over  $E$ ) iff can be written as

$$(2.2) \quad \mathcal{L} = \pi_i^\alpha y_\alpha^i + \lambda,$$

where  $\pi_i^\alpha$  and  $\lambda$  are local functions on  $E$ . To go further, we need the fact that the affine bundle  $p_E: JE \rightarrow E$  has a canonical « linearizing bundle »  $L \rightarrow E$ .

To see this, note that the product bundle  $E \times \mathbb{R} \rightarrow E$  is a vector sub-bundle of  $T^*M \otimes_E TM$  in a canonical way. Hence  $L$  is the vector sub-bundle of  $T^*M \otimes_E TE$  which is projected on  $E \times \mathbb{R}$  by  $id_{T^*M} \otimes_E Tp$ . Moreover, from the canonical injection  $JE \hookrightarrow T^*M \otimes_E TE$  (cf. (1.1)) which takes its values in  $L$ , we get a canonical affine injection  $j: JE \rightarrow L$ . Then the local expression of the unique  $\tilde{\Lambda} \in L^* \otimes_E \bigwedge^m T^*M$  such that  $\Lambda = \tilde{\Lambda} \circ j$  is

$$(2.3) \quad \tilde{\Lambda} = (\pi_i^\alpha \partial_\alpha \otimes dy^i + \lambda \partial_\alpha \otimes dx^\alpha) \otimes \omega.$$

Since there is a canonical injection of

$$L^* \otimes_E \bigwedge^m T^*M \quad \text{into} \quad T^*E \otimes_E \bigwedge^{m-1} T^*M,$$

from  $\Lambda$ , by using the projection  $Tp$ , we get an  $m$ -form  $\Lambda_E$  on  $E$  whose local expression is

$$(2.4) \quad \Lambda_E = \pi_i^\alpha dy^i \wedge \omega_\alpha + \lambda \omega, \quad \omega_\alpha = \partial_\alpha \lrcorner \omega,$$

where  $\lrcorner$  denotes the inner product.

Now let  $\rho$  be a  $p$ -horizontal  $(m-1)$ -form on  $E$  and let

$$(2.5) \quad \rho = \rho^\alpha \omega_\alpha,$$

be its local expression ( $\rho^\alpha$  are local functions on  $E$ ). Then by using the functor  $J$  we get an affine Lagrangian form (of the so called divergence type: cf. [1] [5] and [6]), namely  $\Lambda = \text{div } \rho$ , whose local expression is

$$(2.6) \quad \mathcal{L} = (\partial_i \rho^\alpha) y_\alpha^i + \partial_\alpha \rho^\alpha.$$

In terms of the form  $\Lambda_E$  we have  $\Lambda_E = d\rho$ . Hence  $\Lambda_E$  is closed. Conversely, if  $\Lambda$  is an affine Lagrangian form such that  $d\Lambda_E = 0$ ,  $\Lambda$  is locally a Lagrangian form of the divergence type (as can easily be seen).

Let  $\Lambda$  be a Lagrangian form on  $JE$ . By taking the fiber derivative  $D\Lambda$  (with respect to  $E$ ), followed by an obvious canonical contraction, we get an  $(m-1)$ -form  $\Omega_\Lambda$  on  $JE$  valued on  $V^*E$ , that is

$$(2.7) \quad \Omega_\Lambda : JE \rightarrow V^*E \otimes_E \bigwedge^{m-1} T^*M,$$

whose local expression is

$$(2.8) \quad \Omega_\Lambda = \pi_i^\alpha dy^i \otimes \omega_\alpha, \quad \pi_i^\alpha = \partial_i^\alpha \mathcal{L}.$$

The map  $\Omega_\Lambda$  is the so called Legendre transformation associated to  $\Lambda$  [2]. Note that  $\Lambda$  is affine iff  $D\Omega_\Lambda = 0$ .

The map  $\Omega_\Lambda$  can also be viewed, in a canonical way, as an  $m$ -form on  $JE$ . For this, it is sufficient to observe that the projection  $\mathfrak{g}$ , as defined in (1.2), allows us to construct the canonical injection

$$(2.9) \quad V^*E \otimes_E \bigwedge^{m-1} T^*M \hookrightarrow T^*E \otimes_E \bigwedge^{m-1} T^*E.$$

It is clear that  $\Omega_\Lambda$  is a  $p_E$ -horizontal  $m$ -form on  $JE$  whose local expression is

$$(2.10) \quad \Omega_\Lambda = \pi_i^\alpha \mathfrak{g}^i \wedge \omega_\alpha,$$

as follows from (2.8) and (2.9).

Let  $\varphi : E \rightarrow E$  be an automorphism. Then  $(J\varphi)^*\Lambda$  is again a Lagrangian form and it is easily proved that

$$(2.11) \quad (J\varphi)^*\Omega_\Lambda = \Omega_{(J\varphi)^*\Lambda}.$$

Moreover, let  $u_E$  be a vector field on  $E$ , projectable on  $M$ , and let  $u$  be the 1-jet prolongation of  $u_E$ . Then the infinitesimal version of (2.11) is

$$(2.12) \quad L_u \Omega_\Lambda = \Omega_{L_u \Lambda}.$$

b) Now we consider variations of Lagrangian forms. The basic tool is the subbundle  $\Delta \subset TJE$  [10].

DEFINITION. — Let  $\Lambda$  be a Lagrangian form on  $JE$ . A variation of  $\Lambda$  is a  $p_E$ -horizontal  $m$ -form  $\Theta$  on  $JE$  such that

$$\begin{aligned} i) & \quad \Theta | \Delta = \Lambda | \Delta, \\ ii) & \quad d\Theta | \Delta = 0. \end{aligned}$$

Let  $\{\Lambda\}$  be the set of variations of  $\Lambda$ . Clearly  $\{\Lambda\}$  is an affine space (over  $\mathbb{R}$ ) with vector space the space  $[\Theta]$  of the  $p_E$ -horizontal  $m$ -form  $\Theta$  on  $JE$  such that  $\Theta | \Delta = 0$  and  $d\Theta | \Delta = 0$ . Note that  $\Lambda \in \{\Lambda\}$  iff  $\Omega_\Lambda = 0$ . There is a canonical form  $\Theta_\Lambda \in \{\Lambda\}$ , namely the so called Poincaré-Cartan form

$$(2.13) \quad \Theta_\Lambda = \Lambda + \Omega_\Lambda.$$



The assignment  $\Lambda \mapsto \{ \Lambda \}$  is functorial, i. e. if  $\varphi : E \rightarrow E$  is an automorphism, we find that

$$(2.14) \quad (J\varphi)^* \{ \Lambda \} = \{ (J\varphi)^* \Lambda \}.$$

In particular, from (2.11) and (2.13) it follows that

$$(2.15) \quad (J\varphi)^* \Theta_\Lambda = \Theta_{(J\varphi)^* \Lambda}.$$

If  $u$  is an i. c. t. on JE, we have

$$(2.16) \quad L_u \{ \Lambda \} \subset \{ L_u \Lambda \}.$$

More particularly, if  $u$  is the 1-jet prolongation of a vector field  $u_E$  on E, projectable on M, from (2.12) and (2.13) we get the infinitesimal version of (2.15), namely

$$(2.17) \quad L_u \Theta_\Lambda = \Theta_{L_u \Lambda}.$$

Note that, since  $\Theta_\Lambda = 0$  iff  $\Lambda = 0$  (as is easily seen), from (2.17) it follows that

$$(2.18) \quad L_u \Theta_\Lambda = 0 \Leftrightarrow L_u \Lambda = 0.$$

c) At least in some particular cases, we can say more about the set of variational forms  $\{ \Lambda \}$ . For example, in Classical Analytical Mechanics one is interested to the case in which  $M = \mathbb{R}$ , the absolute time. Another case which occurs often in practice is that in which the fibers of E are 1-dimensional. In both these cases  $[\bar{\Theta}]$  is trivial.

**PROPOSITION.** — Let  $\Lambda$  be a Lagrangian form on JE. Then if  $\dim M = 1$  or also if  $\dim E = m + 1$ , the Poincaré-Cartan form is the unique variational form of  $\Lambda$ .

*Proof.* — Let  $\dim M = 1$  and let  $\bar{\Theta}$  be a  $p_E$ -horizontal 1-form on JE such that  $\bar{\Theta} | \Delta = 0$ . Then locally we must have

$$(2.19) \quad \bar{\Theta} = f_i \vartheta^i,$$

where  $f_i$  are local functions on JE. The result follows now from

$$(2.20) \quad \partial_i^1 \lrcorner d\bar{\Theta} = -f_i dx^1 + (\partial_i^1 f_j) \vartheta^j.$$

The case  $\dim E = m + 1$  is treated in the same way.

Apart from these two cases, in general  $[\bar{\Theta}]$  is not trivial. For example, let  $M = \mathbb{R}^2$  and let  $E = \mathbb{R}^2 \times \mathbb{R}^2$ . Then the 2-form on JE

$$(2.21) \quad \bar{\Theta} = (y_1^1 y_2^2 - y_2^1 y_1^2) dx^1 \wedge dx^2 + \vartheta^1 \wedge dy^2 + dy^1 \wedge \vartheta^2 - dy^1 \wedge dy^2,$$

belongs to  $[\bar{\Theta}]$ .

### 3. VARIATIONAL STRUCTURES

a) The concept of variational form leads us to introduce, in a natural way,  $m$ -forms defined on  $\Delta$ . This is done in the following proposition.

PROPOSITION. — Let  $\Lambda$  be a Lagrangian form on JE. Then there exists a unique  $m$ -form  $\mathcal{E}_\Lambda$  defined on  $\Delta$  and valued on  $V^*E$  such that

$$(3.1) \quad (js)^*(u_0 \lrcorner d\Theta) = - \langle (js)^*\mathcal{E}_\Lambda, \mathcal{V} \circ u_0 \circ js \rangle,$$

where  $\Theta \in \{ \Lambda \}$ ,  $s : M \rightarrow E$  is any section and  $u_0$  is any vector field on JE. The local expression of  $\mathcal{E}_\Lambda$  is

$$(3.2) \quad (\mathcal{E}_\Lambda)_i = [d\pi_i^\alpha \wedge \omega_\alpha - (\partial_i \mathcal{L})\omega] | \Delta.$$

Proof. — The uniqueness of  $\mathcal{E}_\Lambda$  follows easily from (3.1). To prove the existence, we use the well known formula

$$d\Theta(u_0, u_1, \dots, u_m) = \sum_{i=0}^m (-1)^i u_i \cdot \Theta(u_0, \dots, \hat{u}_i, \dots, u_m) + \sum_{0 \leq i < j \leq m} (-1)^{i+j} \Theta([u_i, u_j], u_0, \dots, \hat{u}_i, \dots, \hat{u}_j, \dots, u_m),$$

where  $u_1, \dots, u_m$  are vector fields on JE belonging to  $\Delta$ . Let us note preliminarily that

$$(3.4) \quad d\Theta(\partial_\alpha, u_1, \dots, u_m) = - y_\alpha^i d\Theta(\partial_i, u_1, \dots, u_m),$$

since  $d\Theta | \Delta = 0$  and the vector fields  $w_\alpha = \partial_\alpha + y_\alpha^i \partial_i \in \Delta$ .

By putting  $u_0 = \partial_i^\alpha$ ,  $u_\alpha = w_\alpha$  into (3.9) and by using the properties of  $\Theta$ , it follows that

$$(3.5) \quad \pi_i^\alpha = (-1)^{\alpha-1} \Theta(\partial_i, w_1, \dots, \hat{w}_\alpha, \dots, w_m),$$

because  $[\partial_i^\alpha, w_\beta] = \partial_\beta^\alpha \partial_i$ .

Let now  $s : M \rightarrow E$  be a section and let

$$(3.6) \quad u_\alpha = w_\alpha + (\partial_{\alpha\beta}^2 s^i) \partial_i^\beta \in \Delta, \quad s^i = y^i \circ s.$$

By putting again these vector fields in (3.9), since  $[u_\alpha, u_\beta] = 0$ , we get

$$(3.7) \quad (js)^*(u_0 \lrcorner d\Theta)(\partial_1, \dots, \partial_m) = d\Theta(u_0, u_1, \dots, u_m) \circ js = (js)^* \left[ u_0 \mathcal{L} + \sum_{\alpha=1}^m (-1)^\alpha u_\alpha \cdot \Theta(u_0, w_1, \dots, \hat{w}_\alpha, \dots, w_m) + \sum_{\alpha=1}^m (-1)^\alpha \Theta([u_0, u_\alpha], w_1, \dots, \hat{w}_\alpha, \dots, w_m) \right],$$

where we have used the properties that  $\Theta$  is  $p_E$ -horizontal and that  $\Theta \mid \Delta = \Lambda \mid \Delta$ . By putting  $u_0 = \partial_i$  into (3.7) and by taking into account (3.5), it follows that

$$(3.8) \quad (js)^*(\partial_i \lrcorner d\Theta)(\partial_1, \dots, \partial_m) = -(js)^*[d\pi_i^\alpha \wedge \omega_\alpha - (\partial_i \mathcal{L})\omega](\partial_1, \dots, \partial_m).$$

Finally, if the local expression of  $u_0$  is  $u_0 = u^\alpha \partial_\alpha + u^i \partial_i + u_\alpha^i \partial_i^\alpha$ , since  $d\Theta \mid \Delta = 0$ , we get from (3.4) and (3.8)

$$(3.9) \quad (js)^*(u_0 \lrcorner d\Theta) = -(js)^*[d\pi_i^\alpha \wedge \omega_\alpha - (\partial_i \mathcal{L})\omega](u^i - y_\alpha^i u^\alpha) \circ js.$$

Hence the proof is complete.

The functoriality of the assignment  $\Lambda \rightarrow \mathcal{E}_\Lambda$  can easily be proved. Let  $\varphi : E \rightarrow E$  be an automorphism (over  $\varphi_M : M \rightarrow M$ ). Then by taking the pull-back of both sides of (3.1) by means of  $\varphi_M$  and by using (1.5), we have

$$(3.10) \quad (js^*)^* \{ [(J\varphi)^*u_0] \lrcorner d(J\varphi)^*\Theta \} = - \langle (js^*)^*\varphi^*\mathcal{E}_\Lambda, \vartheta \circ (J\varphi)^*u_0 \circ js^* \rangle,$$

where  $s^* = \varphi^*s$ . By recalling (2.14) and the uniqueness property of  $\mathcal{E}_{(J\varphi)^*\Lambda}$ , as stated before, (3.10) implies that

$$(3.11) \quad \varphi^*\mathcal{E}_\Lambda = \mathcal{E}_{(J\varphi)^*\Lambda}.$$

b) Let  $\Lambda$  be a Lagrangian form on JE. Let us recall that  $\mathcal{E}_\Lambda$  is an  $m$ -form defined only on  $\Delta \subset TJE$  and valued in  $V^*E$ . Then the second order canonical inclusion (1.6) induces the operator

$$(3.12) \quad E_\Lambda : J^2E \rightarrow V^*E \otimes_E \bigwedge^m T^*M.$$

It is characterized by

$$(3.13) \quad E_\Lambda \circ j^2s = (js)^*\mathcal{E}_\Lambda,$$

for any section  $s : M \rightarrow E$ . This second order operator is the Euler-Lagrange operator. Its local expression, as follows from (3.2) and (3.13), is

$$(3.14) \quad E_\Lambda = [(\partial_j^\beta \pi_i^\alpha) y_{\alpha\beta}^j + (\partial_j \pi_i^\alpha) y_\alpha^j + \partial_\alpha \pi_i^\alpha - \partial_i \mathcal{L}] dy^i \otimes \omega.$$

Note that  $E_\Lambda$  is an affine map over JE.

Following the procedure as in (2.9), it is clear that  $E_\Lambda$  can be viewed as an  $(m+1)$ -form on  $J^2E$  whose local expression is

$$(3.15) \quad E_\Lambda = [(\partial_j^\beta \pi_i^\alpha) y_{\alpha\beta}^j + (\partial_j \pi_i^\alpha) y_\alpha^j + \partial_\alpha \pi_i^\alpha - \partial_i \mathcal{L}] dy^i \wedge \omega.$$

The functoriality of  $\Lambda \mapsto E_\Lambda$  can be seen in this way. If  $\varphi : E \rightarrow E$  is an automorphism, by taking the pull-back of both sides of (3.13) by means of  $\varphi^*$ , we get

$$(3.16) \quad (\varphi^*E_\Lambda) \circ j^2s^* = (js^*)^*\mathcal{E}_{(J\varphi)^*\Lambda}, \quad s^* = \varphi^*s,$$

where we have used (3.11). Hence the uniqueness property of  $E_{(J\varphi)^*\Lambda}$  implies that

$$(3.17) \quad \varphi^*E_\Lambda = E_{(J\varphi)^*\Lambda}.$$

The following proposition gives the infinitesimal version of (3.17) in terms of i. c. t.

**PROPOSITION.** — Let  $u_E$  be a vector field on  $E$ , projectable on  $M$ . Then we have

$$(3.18) \quad L_u E_\Lambda = E_{L_u \Lambda}.$$

where with  $u$  on the right and the left side of (3.18) we denote the 1 and 2-jet prolongation of  $u_E$ , respectively.

*Proof.* — We use the formula

$$(3.19) \quad d\Theta_\Lambda = -E_\Lambda + d_V \Omega_\Lambda,$$

where  $d_V$  is the vertical derivation on jet spaces [12]. Formula (3.19) can be proved by a straightforward calculation. The local expression of  $d_V \Omega_\Lambda$  is

$$(3.20) \quad d_V \Omega_\Lambda = (\partial_j \pi_i^\alpha) \vartheta^j \wedge \vartheta^i \wedge \omega_\alpha + (\partial_j^\beta \pi_i^\alpha) \vartheta_j^\beta \wedge \vartheta^i \wedge \omega_\alpha.$$

Since the following commutation relation holds

$$(3.21) \quad L_u d_V \Omega_\Lambda = d_V L_u \Omega_\Lambda,$$

from (3.19) we get

$$(3.22) \quad d\Theta_{L_u \Lambda} = -L_u E_\Lambda + d_V \Omega_{L_u \Lambda},$$

where we have used (2.12) and (2.17). Hence the result follows by comparing (3.22) with

$$(3.23) \quad d\Theta_{L_u \Lambda} = -E_{L_u \Lambda} + d_V \Omega_{L_u \Lambda}.$$

c) There are some useful relations between  $\Lambda$ ,  $E_\Lambda$  and  $\Theta_\Lambda$ . Note that  $\mathcal{E}_\Lambda = 0$  is equivalent to  $E_\Lambda = 0$ , as follows from (3.13). From (3.1) it follows that  $d\Theta_\Lambda = 0$  implies  $E_\Lambda = 0$  (and also that  $\Lambda$  is affine). Also the converse is true. Hence we have

$$(3.24) \quad d\Theta_\Lambda = 0 \Leftrightarrow \Lambda \text{ is affine and } E_\Lambda = 0.$$

Another statement equivalent to (3.24) is:  $\Lambda$  is affine and  $\Lambda_E$  is closed (cfr. (2.4)).

Let us remark that if  $\dim M = 1$  or also if  $\dim E = m + 1$ , then  $E_\Lambda = 0$  implies that  $\Lambda$  is affine. It follows that also  $d\Theta_\Lambda = 0$ . However, apart from these special cases, in general  $E_\Lambda = 0$  does not implies that  $\Lambda$  is affine [6].

d) Given a Lagrangian form  $\Lambda$ , intuitively, the « variational symmetries of  $\Lambda$  » are the morphisms of a category whose objects are the variational forms  $\{\Lambda\}$ . This is made precise by the following definition.

**DEFINITION.** — Let  $\Lambda$  be a Lagrangian form on  $JE$ . Then an automor-

phism  $\varphi : E \rightarrow E$  is said to be a variational symmetry (in short: a v. s.) of  $\Lambda$  iff there is some variational form  $\Theta \in \{ \Lambda \}$  such that

$$(3.25) \quad (J\varphi)^*d\Theta \in d\{ \Lambda \}.$$

Note that by taking the automorphisms  $\varphi : E \rightarrow E$  such that

$$(3.26) \quad (J\varphi)^*\Theta \in \{ \Lambda \},$$

for some  $\Theta \in \{ \Lambda \}$ , we get a subcategory of the previous one. It is clear that if  $\varphi$  is a v. s. of  $\Lambda$ , then  $\varphi$  is a symmetry of the Euler-Lagrange operator  $E_\Lambda$ , i. e.  $\varphi^*E_\Lambda = E_\Lambda$ . This follows from (3.1) in the same way used to prove (3.10).

The existence of a canonical variational form of  $\Lambda$ , namely the Poincaré-Cartan  $\Theta_\Lambda$ , allows us to consider some distinguished v. s. By recalling (2.15) and the equivalence between  $\Theta_\Lambda = 0$  and  $\Lambda = 0$ , it follows that

$$(3.27) \quad \varphi \in \text{Aut} \{ \Theta_\Lambda \} \Leftrightarrow \varphi^*\Lambda = \Lambda.$$

Hence these v. s. are just the symmetries of  $\Lambda$ . Moreover, by recalling (3.24), we get

$$(3.28) \quad \varphi \in \text{Aut} \{ d\Theta_\Lambda \} \Leftrightarrow (J\varphi)^*\Lambda - \Lambda \text{ is affine and } \varphi^*E_\Lambda = E_\Lambda.$$

Note that if  $\dim M = 1$  or also if  $\dim E = m + 1$ , then a symmetry of  $E_\Lambda$  is also an automorphism of  $d\Theta_\Lambda$ .

We finish by considering briefly the infinitesimal symmetries. Let  $u_E$  be a vector field on  $E$  projectable on  $M_1$  and let  $u$  be its 1-jet prolongation. Then  $u_E$  is said to be an infinitesimal variational symmetry (in short: i. v. s.) of a Lagrangian form  $\Lambda$  iff

$$(3.29) \quad L_u d\Theta = 0,$$

for some  $\Theta \in \{ \Lambda \}$ . It is clear that if  $u_E$  is an i. v. s. of  $\Lambda$ , then  $u$  (the 2-jet prolongation of  $u_E$ ) is an infinitesimal symmetry of the Euler-Lagrange operator  $E_\Lambda$ , i. e.  $L_u E_\Lambda = 0$ , as follows from (3.1) and (3.18).

Clearly, we get i. v. s. of  $\Lambda$  by taking those vector fields  $u_E$  on  $E$ , projectable on  $M$ , such that

$$(3.30) \quad L_u \Theta = 0,$$

for some  $\Theta \in \{ \Lambda \}$  (as before  $u$  is the 1-jet prolongation of  $u_E$ ).

The infinitesimal versions of (3.27) and (3.28) are

$$(3.31) \quad L_u \Theta_\Lambda = 0 \Leftrightarrow L_u \Lambda = 0,$$

and

$$(3.32) \quad L_u d\Theta_\Lambda = 0 \Leftrightarrow L_u \Lambda \text{ is affine and } L_u E_\Lambda = 0,$$

respectively.

### 4. CRITICAL SECTIONS

a) In order to exhibit the functorial nature of a variational problem in the most direct way, we start with the following definition.

DEFINITION. — Let  $s : M \rightarrow E$  be a section. A variation of  $s$  is a pair  $(P, (\psi_t)_*s)$  where

i)  $P \subset M$  is an orientable  $m$ -dimensional compact submanifold of  $M$  with boundary (denoted by  $\partial P$ ),

ii)  $\psi_t$  is a one-parameter family of automorphisms  $\psi_t : E \rightarrow E$  (over  $\chi_t : M \rightarrow M$ ) such that

$$(4.1) \quad \psi_t|_{P^{-1}(\partial P)} = id_{P^{-1}(\partial P)} \quad \text{for any } t.$$

The existence of  $\psi_t$  follows easily by considering, for example, vertical fields on  $E$  with compact support. Note that (4.1) implies that

$$(4.2) \quad \chi_t|_{\partial P} = id_{\partial P}, \quad (\psi_t)_*s|_{\partial P} = s|_{\partial P} \quad \text{for any } t.$$

Variations of  $s$  behave in the right way under automorphisms of  $E$ . In fact, let  $\varphi : E \rightarrow E$  be an automorphism (over  $\varphi_M : M \rightarrow M$ ) and let  $(P, (\psi_t)_*s)$  be a variation of  $s$ . Then

$$(4.3) \quad \varphi^*(P, (\psi_t)_*s) = (N, \varphi^*(\psi_t)_*s),$$

where  $N = \varphi_M^{-1}(P)$ , is a variation of  $\varphi^*s$  since

$$(4.4) \quad \varphi^*(\psi_t)_*s = (\varphi_t)_*\varphi^*s, \quad \varphi_t = \varphi^*\psi_t = \varphi_M^{-1} \circ \psi_t \circ \varphi_M.$$

DEFINITION. — Let  $\Lambda$  be a Lagrangian form on  $JE$ . A section  $s : M \rightarrow E$  is said to be critical (with respect to  $\Lambda$ ) iff the function

$$(4.5) \quad t \mapsto \int_{\chi_t(P)} j[(\psi_t)_*s]^*\Lambda$$

is stationary at  $t = 0$  for any variation  $(P, (\psi_t)_*s)$  of  $s$ .

Let  $\varphi : E \rightarrow E$  be an automorphism (over  $\varphi_M : M \rightarrow M$ ). The  $\varphi^*s$  is a critical section of the Lagrangian form  $(J\varphi)^*\Lambda$  iff  $s$  is a critical section of  $\Lambda$ , as follows from (4.4)

$$(4.6) \quad \int_{\chi_t(P)} j[(\psi_t)_*s]^*\Lambda = \int_{\varphi_M^{-1}(\chi_t(P))} \varphi_M^*j[(\psi_t)_*s]^*\Lambda \\ = \int_{\lambda_t(N)} (J\varphi \circ j[\varphi^*(\psi_t)_*s])^*\Lambda = \int_{\lambda_t(N)} j[(\varphi_t)^*\varphi^*s]^*(J\varphi)^*\Lambda,$$

where  $\lambda_t = \varphi_M^*\chi_t$  is the diffeomorphism induced by  $\varphi_t$ .

b) Given a Lagrangian form  $\Lambda$ , critical sections of  $\Lambda$  can be characterized by using the form  $\mathcal{E}_\Lambda$  as follows.

PROPOSITION. — Let  $\Lambda$  be a Lagrangian form on JE. Then a section  $s : M \rightarrow E$  is critical (with respect to  $\Lambda$ ) iff  $(js)^*\mathcal{E}_\Lambda = 0$ .

*Proof.* — The derivative of the function (4.5) at  $t = 0$  is

$$(4.7) \quad D\left(\int_{X_t(P)} j[(\psi_t)_*s]^*\Lambda\right)\Big|_{t=0} = \int_P (js)^*L_u\Lambda,$$

where  $u$  is the 1-jet prolongation of  $u_E$ , the vector field on  $E$  determined by  $\psi_t$ . This result follows by applying the change of variables formula to (4.5), i. e.

$$(4.8) \quad \int_{X_t(P)} j[(\psi_t)_*s]^*\Lambda = \int_P (js)^*(J\psi_t)^*\Lambda.$$

Note that  $u$  is just the vector field on JE determined by the one parameter family  $J\psi_t$ .

By using successively (2.16), the well known identity

$$L_u\Theta = u \lrcorner d\Theta + d(u \lrcorner \Theta)$$

and Stokes' theorem, (4.7) can be written as

$$(4.9) \quad \int_P (js)^*L_u\Lambda = \int_P (js)^*(u \lrcorner d\Theta) + \int_{\partial P} (js)^*(u \lrcorner \Theta),$$

where  $\Theta \in \{\Lambda\}$ . The last integral vanishes since  $\Theta$  is  $p_E$ -horizontal and  $u_E|_{p^{-1}(\partial P)} = 0$ . The result follows now from (3.1).

By choosing  $\Theta \in \{\Lambda\}$ , from (3.1) we get another characterization of critical sections, namely a section  $s$  is critical iff

$$(4.10) \quad (js)^*(u \lrcorner d\Theta) = 0,$$

for any vector field  $u$  on JE (or also for any i. c. t.  $u$  projectable over  $E$ ). In terms of the Euler-Lagrange operator  $E_\Lambda$ , the condition for  $s$  to be critical is the well known one, namely  $E_\Lambda \circ j^2s = 0$ , as follows from (3.13).

## REFERENCES

- [1] R. COURANT and D. HILBERT, *Methods of Mathematical Physics*, t. 1, New York and London, 1953.
- [2] P. L. GARCIA, The Poincaré-Cartan Invariant in the Calculus of Variations, *Symposia Mathematica*, t. 14, 1974, p. 219-246.
- [3] H. GOLDSCHMIDT and S. STERNBERG, The Hamilton-Cartan Formalism in the Calculus of Variations, *Ann. Inst. Fourier, Grenoble*, t. 23 (1), 1973, p. 203-267.
- [4] R. HERMANN, *Geometry, Physics and Systems*, M. Dekker, Inc., New York, 1973.
- [5] E. L. HILL, Hamilton's Principle and the Conservation Theorems of Mathematical Physics, *Reviews of Modern Physics*, t. 20 (3), 1951, p. 253-260.
- [6] D. KRUPKA, Some Geometric Aspects of Variational Problems in Fibered Manifolds, *Folia Fac. Scie. Nat. Univ. Purkyniane Brunensis*, t. 14, Opus 10, 1973, p. 1-65.

- [7] B. A. KUPERSHMIT, Geometry of Jet Bundles and the Structure of Lagrangian and Hamiltonian Formalisms, *Lecture Notes in Mathematics*, t. **775**, 1979, p. 162-218.
- [8] L. MANGIAROTTI and M. MODUGNO, New Operators on Jet Spaces, to appear.
- [9] R. OUZILOU, Expression Symplectique des Problèmes Variationnels, *Symposia Mathematica*, t. **14**, 1974, p. 85-98.
- [10] J. ŚNIATYCKI, On the Geometric Structure of Classical Field Theory in Lagrangian Formulation, *Proc. Camb. Phil. Soc.*, t. **68**, 1970, p. 475-484.
- [11] A. TRAUTMANN, Noether Equations and Conservation Laws, *Comm. Math. Phys.*, t. **6**, 1967, p. 248-261.
- [12] W. M. TULCZYJEW, The Euler-Lagrange Resolution, *Lecture Notes in Mathematics*, t. **836**, 1980, p. 22-48.

(Manuscrit reçu le 26 mars 1982)